5

10

15

20

## What Is Claimed Is:

- 1. A method for inhibiting angiogenesis in a tissue comprising administering to said tissue a composition comprising an angiogenesis-inhibiting amount of an  $\alpha_{\nu}\beta_{3}$  antagonist.
- 2. The method of claim 1 wherein said  $\alpha_{\nu}\beta_{3}$  antagonist inhibits binding of fibringen to  $\alpha_{\nu}\beta_{3}$  but does not substantially inhibit binding of fibringen to  $\alpha_{\text{IIb}}\beta_{3}$ .
- 3. The method of claim 1 wherein said  $\alpha_v \beta_3$  antagonist is a monoclonal antibody immunospecific for  $\alpha_v \beta_3$ .
- 4. The method of claim 3 wherein said monoclonal antibody has the immunoreaction characteristics of the monoclonal antibody LM609.
- 5. The method of claim/1 wherein said  $\alpha_{\rm v}\beta_3$  antagonist is an RGD-containing polypeptide.
- 6. The method of claim 5 wherein said polypeptide is selected from the group consisting of c-(GrGDFV), c-(RGDFV), c-(RADFV), c-(RGDFV) and YTAECKPQVTRGDVF.

  and a salt thereof.
- 7. The method of claim 6 wherein said salt is hydrochloride or trifluoroacetate.
- 8. The method of claim 1 wherein said tissue is inflamed and said angiogenesis is inflamed tissue angiogenesis
- 9. The method of claim 8 wherein said tissue is arthritic.
- 10. The method of claim 9 wherein said arthritic tissue is present in a mammal with rheumatoid arthritis.
- 11. The method of claim 1 wherein said tissue is the retinal tissue of a patient with diabetic retinopathy and said angiogenesis is retinal

75

25

30

35

angiogenesis.

- 12. The method of claim 1 wherein said tissue is a hemangioma.
- 13. The method of claim 1 wherein said tissue is a solid tumor or a solid tumor metastasis and said angiogenesis is tumor angiogenesis.
- 14. The method of claim 1 wherein said angiogenesis-inhibiting amount is from about 2 uM to 5 mM.
- 15. The method of claim in wherein said administering comprises intravenous, transdermal, intrasynovial, intramuscular, or oral administration.
- 16. The method of claim 1 wherein said administering is conducted in conjunction with chemotherapy.

15

10

5

Kg (

